

BRITISH GEOLOGICAL SURVEY

King Edward Point

Observatory

Monthly Magnetic Bulletin

December 2018

18/12/KE



**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

KING EDWARD POINT OBSERVATORY MAGNETIC DATA

1. Introduction

King Edward Point observatory was installed by the British Geological Survey (BGS), with the permission of the Government of South Georgia and the South Sandwich Islands (GSGSSI), in collaboration with British Antarctic Survey (BAS), and became operational in February 2011.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team
British Geological Survey
Lyell Centre, Heriot Watt University
Research Avenue South
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
E-mail: enquiries@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

King Edward Point Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site adjacent to the Government of South Georgia and the South Sandwich Islands (GSGSSI) research station at King Edward Point, Cumberland East Bay, South Georgia. The observatory co-ordinates are:

Geographic: 54° 16' 55.2" S 323° 30' 25.2" E
Geomagnetic: 47° 38' 60" S 030° 14' 24" E
Height above mean sea level: 7 m

The geographical coordinates are measured by a handheld GPS device, which uses WGS84 as the reference coordinate system. The height above MSL is determined from the best available contour maps. The geomagnetic co-ordinates are approximations, calculated using the 12th generation International Geomagnetic Reference Field (IGRF) at epoch 2018.5. On-line access to models (including IGRF), charts and navigational data are available at

http://www.geomag.bgs.ac.uk/data_service/models_compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in February 2011. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at www.geomag.bgs.ac.uk/data_service/home

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. One set of absolute measurements of the field are made manually once per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically

quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at King Edward Point magnetic observatory, South Georgia operated by the British Geological Survey in co-operation with the British Antarctic Survey and the Government of South Georgia and the South Sandwich Islands (GSGSSI).'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff

KING EDWARD POINT OBSERVATORY


ABSOLUTE OBSERVATIONS

Date	Day Number	Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity		Observer
		Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	
03-Dec-18	337	16:53	-6.9829	-7.0233	17:13	-56.0106	12.4	27696.1	15483.2	15997.3	-22964.0	-23200.0	AC
03-Dec-18	337	17:29	-7.0129	-7.0267	17:47	-55.9831	12.4	27703.1	15498.1	15998.2	-22962.3	-23199.3	AC
19-Dec-18	353	11:00	-1.3350	999.9999	11:15	-56.0742	12.4	27720.1	15471.1	99999.9	-23001.1	99999.9	AC
19-Dec-18	353	11:26	-7.1547	-7.0267	11:39	-55.9335	12.4	27720.6	15527.8	15997.2	-22963.4	-23199.7	AC

Note:

1: 999.9999 and 99999.9 denote a flagged value


King Edward Point 2018




King Edward Point

December

2018






Date: 01-12-2018


King Edward Point

Day number: 335



Date: 02-12-2018


Day number: 336



Date: 03-12-2018


King Edward Point

Day number: 337



Date: 04-12-2018


Day number: 338



Date: 05-12-2018


King Edward Point

Day number: 339



Date: 06-12-2018


Day number: 340



Date: 07-12-2018


King Edward Point

Day number: 341



Date: 08-12-2018


Day number: 342



Date: 09-12-2018


King Edward Point

Day number: 343



Date: 10-12-2018


Day number: 344



Date: 11-12-2018


King Edward Point

Day number: 345



Date: 12-12-2018


Day number: 346



Date: 13-12-2018


King Edward Point

Day number: 347



Date: 14-12-2018


Day number: 348



Date: 15-12-2018


King Edward Point

Day number: 349



Date: 16-12-2018


Day number: 350



Date: 17-12-2018


King Edward Point

Day number: 351



Date: 18-12-2018


Day number: 352



Date: 19-12-2018


King Edward Point

Day number: 353



Date: 20-12-2018


Day number: 354



Date: 21-12-2018


King Edward Point

Day number: 355



Date: 22-12-2018


Day number: 356



Date: 23-12-2018


King Edward Point

Day number: 357



Date: 24-12-2018


Day number: 358



Date: 25-12-2018


King Edward Point

Day number: 359



Date: 26-12-2018


Day number: 360



Date: 27-12-2018


King Edward Point

Day number: 361



Date: 28-12-2018


Day number: 362



Date: 29-12-2018


King Edward Point

Day number: 363



Date: 30-12-2018


Day number: 364




Date: 31-12-2018

King Edward Point

Day number: 365




King Edward Point Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

King Edward Point Observatory: Horizontal Intensity (nT)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

King Edward Point Observatory: Vertical Intensity (nT)



King Edward Point Observatory 2018



Monthly Mean Values for King Edward Point Observatory 2018

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	-7° 9.5′	15571 nT	-55° 53.6′	15449 nT	-1940 nT	-22992 nT	27769 nT
February	-7° 8.9′	15559 nT	-55° 54.7′	15438 nT	-1936 nT	-22990 nT	27761 nT
March	-7° 9.0′	15548 nT	-55° 55.9′	15427 nT	-1935 nT	-22992 nT	27755 nT
April	-7° 8.8′	15543 nT	-55° 56.2′	15422 nT	-1934 nT	-22989 nT	27750 nT
May	-7° 8.8′	15538 nT	-55° 56.6′	15417 nT	-1933 nT	-22987 nT	27746 nT
June	-7° 9.0′	15534 nT	-55° 56.8′	15413 nT	-1933 nT	-22983 nT	27740 nT
July	-7° 8.7′	15533 nT	-55° 56.6′	15412 nT	-1932 nT	-22979 nT	27736 nT
August	-7° 8.6′	15522 nT	-55° 57.5′	15402 nT	-1930 nT	-22977 nT	27729 nT
September	-7° 8.2′	15516 nT	-55° 58.2′	15396 nT	-1928 nT	-22978 nT	27726 nT
October	-7° 7.9′	15518 nT	-55° 57.7′	15398 nT	-1926 nT	-22973 nT	27723 nT
November	-7° 7.6′	15517 nT	-55° 57.4′	15398 nT	-1925 nT	-22968 nT	27718 nT
December	-7° 7.3′	15517 nT	-55° 57.2′	15398 nT	-1924 nT	-22965 nT	27716 nT

Note

i. The values shown here are provisional.